


A Step-by-Step Guide for the Production of Recombinant Fluorescent TAT-HA-Tagged Proteins and their Transduction into Mammalian Cells

Christer Abou Anny,¹ Sébastien Nouaille,² Régis Fauré,² Céline Schulz,¹ Corentin Spriet,^{1,3} Isabelle Huvent,¹ Christophe Biot,¹ and Tony Lefebvre^{1,4} 

¹Université de Lille, CNRS, UMR 8576 - UGSF, Lille, France

²TBI, Université de Toulouse, CNRS, INRAE, INSA, Toulouse, France

³Université de Lille, CNRS, Inserm, CHU Lille, Institut Pasteur de Lille, US 41 - UAR 2014 - PLBS, F-59000, Lille, France

⁴Corresponding author: tony.lefebvre@univ-lille.fr

Published in the Molecular Biology section

Investigating the function of target proteins for functional prospection or therapeutic applications typically requires the production and purification of recombinant proteins. The fusion of these proteins with tag peptides and fluorescently derived proteins allows the monitoring of candidate proteins using SDS-PAGE coupled with western blotting and fluorescent microscopy, respectively. However, protein engineering poses a significant challenge for many researchers. In this protocol, we describe step-by-step the engineering of a recombinant protein with various tags: TAT-HA (trans-activator of transduction-hemagglutinin), 6×His and EGFP (enhanced green fluorescent protein) or mCherry. Fusion proteins are produced in *E. coli* BL21(DE3) cells and purified by immobilized metal affinity chromatography (IMAC) using a Ni-nitrilotriacetic acid (NTA) column. Then, tagged recombinant proteins are introduced into cultured animal cells by using the penetrating peptide TAT-HA. Here, we present a thorough protocol providing a detailed guide encompassing every critical step from plasmid DNA molecular assembly to protein expression and subsequent purification and outlines the conditions necessary for protein transduction technology into animal cells in a comprehensive manner. We believe that this protocol will be a valuable resource for researchers seeking an exhaustive, step-by-step guide for the successful production and purification of recombinant proteins and their entry by transduction within living cells. © 2024 The Authors. Current Protocols published by Wiley Periodicals LLC.

Basic Protocol 1: DNA cloning, molecular assembly strategies, and protein production

Basic Protocol 2: Protein purification

Basic Protocol 3: Protein transduction in mammalian cells

Keywords: DNA molecular assembly • fluorescent protein • fusion protein • Ni-NTA-based protein purification • protein transduction • recombinant protein production • tagged protein

How to cite this article:

Anny, C. A., Nouaille, S., Fauré, R., Schulz, C., Spriet, C., Huvent, I., Biot, C., & Lefebvre, T. (2024). A step-by-step guide for the production of recombinant fluorescent TAT-HA-Tagged proteins and their transduction into mammalian cells. *Current Protocols*, 4, e1016. doi: 10.1002/cpz1.1016

INTRODUCTION

Over the years, the interest in innovative approaches for the expression, purification, and characterization of recombinant proteins has increased. It empowers scientists to create, engineer, and produce proteins with diverse functions, characteristics, and applications (Assenberg et al., 2013). Thus, production of recombinant proteins represents a cornerstone for various applications in drug discovery (Andersen & Krummen, 2002; Overton, 2014), therapeutic proteins (Burnett & Burnett, 2020; Wurm, 2004) or vaccine production (Farzaneh et al., 2017; Joung et al., 2016). Recombinant DNA encoding proteins are typically generated through the use of recombinant DNA technology from historical methods (e.g., amplifications and use of restriction enzymes) to recently developed sophisticated technologies (e.g., Gibson assembly method for DNA cloning) (Bordat et al., 2015; Casini et al., 2015; Cohen, 2013; Ferro et al., 2019; Gibson et al., 2009; Jang & Magnuson, 2013; Kostylev et al., 2015; Li et al., 2019; Li et al., 2020; Li et al., 2018; Norris et al., 2015; Roberts, 2019; Rudenko & Barnes, 2018; Sands & Brent, 2016; Thomas et al., 2015; Wang et al., 2015). One of the most significant breakthroughs in recombinant protein technology has been the development of expression systems for the efficient production of complex proteins by selecting an appropriate method in microbial hosts (Brondyk, 2009), such as *Escherichia coli* (Baneyx, 1999; Baneyx & Mujacic, 2004; Chen, 2012; Gopal & Kumar, 2013; Rosano & Ceccarelli, 2014; Sørensen & Mortensen, 2005), yeast (Cregg et al., 2000; Mattanovich et al., 2012; Porro et al., 2005), *Streptomyces spp.* (Anné et al., 2012), mammalian cells (Andersen & Krummen, 2002; Barnes et al., 2003; Hopkins et al., 2012; Jäger et al., 2015; Sunley & Butler, 2010) or plants (Anné et al., 2012; Newman et al., 2011; Nienhaus & Nienhaus, 2014; Noguchi & Matsumoto, 2006). These systems provide researchers with the ability to produce proteins that may be challenging to obtain by fastidious protein purification from various biological sources (Brondyk, 2009). Among these recombinant proteins, fluorescent proteins have proven to be invaluable tools for monitoring and visualizing cellular processes in real-time (Ai et al., 2014; Nienhaus & Nienhaus, 2014; Wang et al., 2008; Wiedenmann et al., 2009). Fluorescent proteins are derived from naturally occurring proteins with intrinsic fluorescence, such as the Green Fluorescent Protein (GFP) originally isolated from *Aequorea victoria* (Jakobs et al., 2000; McRae et al., 2005; Nolte et al., 2001) or the red fluorescent protein mCherry, derived from DsRed isolated from *Discosoma* sea anemones (Shen et al., 2017). These color-differentiated fluorescent proteins can be used together to perform multicolor imaging experiments in molecular and cellular biology research in order to label specific cellular structures, organelles, or proteins for visualization and tracking within living cells (Helmuth et al., 2009; Kirchhausen, 2009; McDonald et al., 2002; Newman et al., 2011; Rizzuto et al., 1995; Salomonsson et al., 2012; Subach et al., 2009; Yin et al., 2013).

This protocol is designed to provide a step-by-step guide to successfully express recombinant fluorescent proteins in bacteria and subsequently visualize them in animal cells (Fig. 1). The successful execution of this protocol involves several steps, starting with the design and construction of recombinant DNA vectors using the Gibson assembly method to express the target protein fused with the HIV-1 trans-activator peptide called TAT-tag




Figure 1 Schematic overview of protocols workflow. This Current Protocol article is divided into three distinct Basic Protocols aimed at cloning by molecular assembly a POI with various tags and producing it in *E. coli* cells (Basic Protocol 1, panel A), to purify this protein (Basic Protocol 2, panel B), and transducing it in eukaryotic animal cells for further investigations (Basic Protocol 3, panel C). POI, Protein of Interest.

(Trans-Activator of Transduction), a fluorescent protein, and a TEV protease recognition sequence to eliminate the downstream 6×His-tag. This method is used for the purpose of gene cloning and expression, through constructing plasmids for protein expression. These assembled vectors are then transformed into a suitable host organism, typically *E. coli* BL21(DE3), to produce recombinant fluorescent proteins. Recombinant protein production is carried out through transcription induction by L-arabinose thanks to the P_{BAD} bacterial promoter (Greenblatt & Schleif, 1971; Schleif, 2000; Schleif et al., 1973). The protein purification is performed via immobilized metal affinity chromatography (IMAC) thanks to the 6×His-tag (Block et al., 2009; Gaberc-Porekar & Menart, 2001; Porath, 1992; Sun et al., 2005). The TAT-tag, due to its transduction property, enables the fused protein to enter animal cells (Becker-Hapak et al., 2001; Eguchi et al., 2001; Fittipaldi & Giacca, 2005; Ford et al., 2001; Gump & Dowdy, 2007; Gump et al., 2010; Guo et al., 2012; Kabouridis, 2003; Kaplan et al., 2005; Leifert et al., 2002; MacKay & Szoka, 2003; Noguchi & Matsumoto, 2006; Pawson & Nash, 2000; Schwarze et al., 2000; Shokolenko et al., 2005; Wadia & Dowdy, 2002; Zhang et al., 2012). Once inside the living cells, the recombinant protein can be monitored and tracked according to its localization and behavior by its fluorescent signal.

The production of proteins in bacteria and their delivery to animal cells by a penetration peptide has the advantage of working with well-defined quantities of proteins, obtaining a homogeneous distribution of the latter throughout the cell population and to be a vector-free method. The transient transfection of vectors, e.g., plasmids, makes it impossible to control the quantity of protein expressed within the cell and very often leads to heterogeneous expression within the cell population, as well as being cell type dependent.

We offer the reader, with a range of readily accessible techniques and devices found in research laboratories or with a simple Ni-NTA column, the opportunity to produce

fluorescent proteins using three distinct but complementary protocols. Basic Protocol 1 details, thanks to Gibson-type molecular assembly, the construction of custom plasmid vectors making it possible to express a specific protein of interest (POI) in phase with any fluorescent protein and various tags, and to produce it in *E. coli* bacteria. Basic Protocol 2 presents the purification of the POI by taking advantage of a poly-His-tag by using a preparative chromatography system with a fast protein liquid chromatography (FPLC) Ni-nitrilotriacetic acid (NTA) agarose column, and its quality control by SDS-PAGE coupled to western blotting. Lastly, Basic Protocol 3 exposes a very simple experimental strategy for protein transduction into animal cells using a cell penetration peptide. We use direct fluorescence microscopy to avoid long, complex, and often costly indirect immunofluorescence steps.

**BASIC
PROTOCOL 1**

**DNA CLONING, MOLECULAR ASSEMBLY STRATEGIES, AND PROTEIN
PRODUCTION**

In this protocol, we describe a procedure to construct a specific DNA sequence encompassing the different blocks needed for expression of the final protein containing the protein of interest (POI) moiety (Fig. 1A and Fig. 2). For that purpose, the choice of the gene




Figure 2 DNA assembly towards the overall plasmid constructions. **(A)** Use of P1-P6 primers to generate the pBAD-TAT-HA-EGFP-TEV-6×His final construction. **(B)** Use of P8-P10 primers to generate the pBAD-TAT-HA-POI-EGFP-TEV-6×His final construction. **(C)** Use of P11-P14 primers to generate the pBAD-TAT-HA-POI-mCherry-TEV-6×His final construction. P1-P14 represent the different primers listed in Table 3 needed to generate the different PCR products that are then PCR-amplified to step-by-step reach the final constructions. For more details see Tables 1–7. The color code used is the same as in Table 3. EGFP, Enhanced Green Fluorescent Protein, LacZ, β-galactosidase encoding gene; Ori, Origin of replication; pBAD, promoter of *araB*, *araA* and *araD*; POI, Protein Of Interest; TAT-HA, Trans-Activator of Transduction-HemAgglutinin TEV, recognition sequence by the Tobacco Etch Virus protease; vCMV, CytoMegalovirus vector.

Table 1 Characteristics of the PCR Components

Plasmid DNA source	Primer pair	Amplicon size (bp)
pBAD-LacZ-X-6×His	P1-P2	4047
pCDNA3-Y-TAT-HA	P3-P4	122
vCMV-EGFP-POI	P5-P6	745
pBAD-TAT-HA-EGFP-TEV-6×His	P7-P8	4873
vCMV-EGFP-POI	P9-P10	2360
pBAD-TAT-HA-POI-EGFP-TEV-6×His	P11-P12	6479
pCMV-mCherry	P13-P14	745

Table 2 Bacterial Strains and Plasmids Used in this Study

Strain or Plasmid	Characteristics	Source
Strain		
Competent <i>E. coli</i> (High Efficiency)	DH5-alpha	New England BioLabs
Competent <i>E. coli</i> (High Efficiency)	BL21(DE3)	New England BioLabs
Plasmid		
pBAD-LacZ-6×His	<i>E. coli</i> expression vector, amp ^r	Life Science
pCDNA3-Y-TAT-HA	<i>E. coli</i> expression vector, amp ^r	UGSF, Lille (France)
vCMVp-EGFP-POI	<i>E. coli</i> expression vector, kan ^r	e-Zyvec, Loos (France)
pBAD-TAT-HA-EGFP-TEV-6×His	<i>E. coli</i> expression vector, amp ^r	This study
pBAD-TAT-HA-POI-EGFP-TEV-6×His	<i>E. coli</i> expression vector, amp ^r	This study
pCMV-mCherry	<i>E. coli</i> expression vector, kan ^r	UGSF, Lille (France)

encoding the POI, the plasmid vector and its features highly depend on specific needs of the experiment, including the type of host organism (e.g., *E. coli*) and the size of the DNA to be inserted. For the expression in *E. coli*, the desired level of gene expression is dependent on the choice of the appropriate promoter (e.g., P_{BAD} promoter, used here). These choices greatly influence the success of gene expression studies (for detailed information see Tables 1–3). The plasmids undergo construction through Gibson molecular assembly of purified PCR fragments. The advantage with this approach is that it is possible, with a whole set of constructions either available in the laboratory or obtained elsewhere, to recover a fragment of interest (comparable to a block or a brick from a game assembly) and nesting it with other fragments to custom manufactured plasmids expressing the POI in phase with any desired tag or fluorescent protein.

Materials

Plasmids, or other source of DNA insert (see Table 2 for detailed information)
 Custom primers (see Table 3 for primers used in this study)
 Q5 high-fidelity DNA polymerase, 2000 U/ml (New England BioLabs, cat. no. M0491S)
 Q5 reaction buffer, 5× (New England BioLabs, cat. no. B9027S)
 Deoxynucleotide (dNTP) solution mix, 10 mM (New England BioLabs, cat. no. N0447S)
 Ultrapure DNase-free H₂O
 Agarose powder
 Tris-acetate-EDTA (TAE) buffer, 0.5× (Euromedex)
 Gel loading buffer (New England Biolabs, cat. no. B7025)

Anny et al.

5 of 26

Table 3 Primers Used in this Study^a

Primer	Sequence (5' → 3')	Construct
P1-TEV-6×His	GAAAACCTGTACTTCCAGGGTAATAGCGCCGTCGACCATCAT	pBAD-TAT-HA-EGFP-TEV-6×His
P2-TAT-HA-pBAD	AGCCAAGCTTTGGATCCATGGTTAATTCCTCCTGT	
P3-pBAD-TAT-HA	CATGGATCCAAGCTTGGCTACGGCCGCAAGAAA	
P4-EGFP-TAT-HA	CCTTGCTCACGGCCATGGAGCCAGCATAGTCTG	
P5-EGFP-TAT-HA	CTCCATGGCCGTGAGCAAGGGCGAGGAGCTGTTC	
P6-TEV-EGFP	ACCCTGGAAGTACAGGTTTTCCTTGACAGCTCGTCCATGCCGAG	
P7-POI-EGFP	NNNNNNNNNNGTGAGCAAGGGCGAGGAGCTGTTC	pBAD-TAT-HA-POI-EGFP-TEV-6×His
P8-POI-TAT-HA	NNNNNNNNNNGGCCATGGAGCCAGCATAGTCTG	
P9-TAT-HA-POI	CTCCATGGCCNNNNNNNNNNNNNNNNNNNNNNNN	
P10-EGFP-POI	CCTTGCTCACNNNNNNNNNNNNNNNNNNNNNNNN	
hP11-POI Rv	CAGGTCAGTATCAAACCAGGCCAGC	pBAD-TAT-HA-POI-mcherry-TEV-6×His
P12-5'TEV	GAAAACCTGTACTTCCAGGGTAATAGCG	
P13-mcherry	CCTGGTTTGATACTGACCTGGTGAGCAAGGGCGAGGAGGATAACAT	
P14-mcherry	CCCTGGAAGTACAGGTTTTCCTTGACAGCTCGTCCATGCCGCC	

^a For clarity the color code used for sequences is the same as that used in Figure 2 for fragments.

- 1 kb Plus DNA ladder, 1000 µg/ml (New England BioLabs, cat. no. N3200S)
- 100 bp DNA ladder, 500 µg/ml (New England BioLabs, cat. no. N3231S)
- Ethidium bromide (Thermo Fisher Scientific, cat. no. 15585011)
- QIAquick gel extraction kit (Qiagen, cat. no. 28706)
- NEBuilder HiFi DNA assembly master mix (New England BioLabs, cat. no. E2621S)
- Competent 5-alpha *E. coli* (New England BioLabs, cat. no. C2987H)
- SOC medium (New England BioLabs, cat. no. B9020S)
- LB agar plates: LB medium plus 15 g/L agar
- LB medium: 10 g/L tryptone, 10 g/L NaCl, and 5 g/L yeast extract supplemented with the appropriate antibiotics, e.g., 100 mg/ml ampicillin (stock solution) diluted at 1/1000
- QIAprep spin miniprep kit (Qiagen, cat. no. 27104)
- Competent BL21(DE3) *E. coli* (New England BioLabs, cat. no. C2527H)
- L-arabinose (Sigma-Aldrich, cat. no. A3256)
- Complete EDTA-free protease inhibitor cocktail tablets (Sigma-Aldrich, cat. no. 11873580001)
- Buffer A (see recipe)
- DNase I (3000 U/mg) 20,000 U (Euromedex, cat. no. 1307)
- 0.2-ml PCR tubes (Dutsher, cat. no. 010207)
- Benchtop centrifuge
- Thermal cycler (Eppendorf Mastercycler)
- Microwave oven or hotplate
- Agarose gel electrophoresis system
- E-Box gel documentation imaging system (Vilber Lourmat, France)
- Benchtop UV transilluminator machine
- Scalpel
- NanoDrop ND-1000 spectrophotometer
- Petri dishes (TRP, cat. no. 067002)
- 1.5-ml microcentrifuge tubes (Thermo Fisher Scientific, cat. no. 69715)
- Ice
- Heat block (Eppendorf ThermoMixer) or water bath
- Shaking incubator
- Sterile toothpicks

Table 4 PCR Components and Procedure for Q5 High-Fidelity DNA Polymerase

Composition of PCR mixture			
Component	Stock solution	Volume (μl)	Final concentration
Q5 reaction buffer	5×	5 μl	1×
dNTP mix	10 mM each	0.5 μl	0.2 mM each
Primers (10 μM)	10 μM	1.25 μl for each primer	0.5 μM each
Plasmid matrix	Variable	Variable	~ 4 pg/μl
Q5 polymerase	2 U/μl	0.3 μl	0.024 U/μl
Ultrapure water		To 25 μl	
PCR Thermocycling conditions			
Step	Temperature (°C)	Time	Cycle
Denaturation	98	30 s	1
Denaturation	98	10 s	
Annealing	60	20 s	×30
Elongation	72	30 s/kb	
Elongation	72	3 min	1
Cooling	14	∞	-

1-L glass Erlenmeyer flasks

Centrifuge Beckman-Coulter Avanti J-26S XP (with a rotor adapted for 1-L volumes)

50-ml sterile conical tubes (Falcon, cat. no. 352070)

Sonicator

Centrifuge, rotor 19776H, angle rotor, for six 50-ml culture tubes

0.45-μm filter (Dutscher, cat. no. 033942)

CAUTION: Ethidium bromide is a putative carcinogen compound (the dye intercalates between the stacked bases of DNA fragments) and must be handled with gloves. Do not use ethidium bromide powder to avoid irritation by inhalation and never use a microwave oven to heat solutions containing ethidium bromide. It is preferable to use the less toxic SYBR safe stain gel to visualize DNA fragments in agarose gels. When using the transilluminator to visualize DNA bands, it is essential to protect yourself from aggressive UV rays with plastic glasses, gloves and sleeves that must cover the integrity of the arms.

Amplification of the various PCR products

1. In PCR tubes, mix the Q5 reaction buffer (adjusted with DNase-free ultrapure water) with the DNA templates, primers (designed as P1 to P14 in Table 1 and Fig. 2), dNTP and the Q5 high-fidelity DNA polymerase at desired concentrations (Table 4). Mix and pulse-centrifuge to ensure the total liquid is located at the bottom of the tube.

Primers P1 and P2 serve to recover the P_{BAD} -LacZ-6×His part; primers P3 and P4, the TAT-HA penetrating peptide encoding part; primers P5 and P6, the EGFP encoding part; primers P7 and P8 serve to recover the $pBAD$ -TAT-HA-EGFP-TEV-6×His by adding 15 to 20 bp of the POI between the TAT-HA and EGFP sequences; primers P9 and P10, the POI encoding part; primers P11 and P12 serve to recover the $pBAD$ -TAT-HA-POI-TEV-6×His by excluding the EGFP part; and primers P13 and P14 serve to recover the mCherry encoding part (Fig. 2).

2. Place the PCR tubes in a thermal cycler and apply the corresponding PCR amplification program (Table 4).




Figure 3 PCR electrophoresis gels. The PCR products of the different fragments shown schematically in Figure 2 were analyzed by electrophoresis on agarose gel. At the top of each gel are indicated the primer pairs following the same color code that used in Figure 2. **(A)** pBAD-TEV-6×His, POI and EGFP fragments. **(B)** TAT-HA fragment. **(C)** POI and pBAD-EGFP-TEV-6×His fragments. **(D)** pBAD-TAT-HA-POI-TEV-6×His and mCherry fragments. The size markers are indicated at the left of each gel: 1 kb Plus DNA ladder for panels **(A)**, **(C)** and **(D)**; 100 bp ladder for panel **(B)**.

3. Prepare the gel by mixing agarose at a concentration of 1% (w/v) in 0.5× TAE. Heat the solution in a microwave oven or on a hotplate until the agarose has dissolved.
4. Pour the gel in the electrophoresis casting tray and let it solidify. Remove the comb and put the gel in the electrophoresis chamber containing 0.5× TAE as migration buffer.
5. Mix the 5 µl PCR reaction to 2 µl gel loading buffer, load on the gel along with 5 µl DNA size marker to monitor DNA size after migration. Migrate at 100 V for 30 min.
6. After migration, incubate the gel for 10 min in water containing 0.5 µg/ml ethidium bromide. Wash the excess of ethidium bromide by incubation in water for 5 min.
7. Visualize the PCR products (Fig. 3) by exposition of the gel to a UV transilluminator, such as the E-box gel documentation imaging system.
8. To purify the amplicon of interest, cut out the UV-detected DNA bands using a scalpel blade under a UV transilluminator instrument.

Table 5 Plasmid DNA Assembly Conditions

Components	Stock solution	Volume	Final concentration
Purified PCR	Variable	Variable	5-15 ng/μl
T4 polynucleotide kinase PNK	10 U/μl	1 μl	0.5 U/μl
T4 DNA ligase Buffer with ATP	10× with 10 mM	1 μl	0.5× with 0.5 mM
T4 DNA ligase	400 U/μl	1 μl	20 U/μl
Ultrapure water		To 20 μl	

Table 6 Gibson Method Plasmid DNA Assembly

Component	Volume (μl)
Purified PCR fragments	x μl
Enzymes, 2× Hi-Fi DNA assembly master mix	5 μl
Water	Up to 10 μl
DNA assembly thermocycling conditions	
50°C for 1 hr	
14°C/∞	

9. Weigh the DNA containing the gel slice and add 3 volumes (v/w) of QG buffer. Purify by using QIAquick gel extraction kit following the manufacturer's instructions. Elute with 50 μl elution buffer. Quantify the DNA using a NanoDrop ND-1000 spectrophotometer.

DNA molecular assembly

10. Mix each purified PCR fragment, i.e., TAT-HA, POI, pBAD-TEV-6×His, and EGFP or mCherry for the construction of pBAD-TAT-HA-POI-EGFP-TEV-6×His and pBAD-TAT-HA-POI-mCherry-TEV-6×His, respectively (Fig. 2), with NEBuilder HiFi DNA assembly 2× master mix at a ratio (1:1) (Tables 5 and 6). Complement with water to a final volume of 10 μl. Incubate the mixture at 50°C for 1 hr in a thermocycler.
11. Use 5 μl assembly reaction for bacterial transformation in *E. coli* 5-alpha chemically competent cells (Froger & Hall, 2007). Mix DNA to 50 μl cells in 1.5-ml microcentrifuge tubes and mix gently by pipetting up and down or flicking the tube 4 to 5 times (do not vortex).
12. After 20 min on ice, perform heat shock using a heat block or water bath set to 42°C for 30 s and immediately transfer cells on ice for 3 min.
13. Add 450 μl SOC medium and incubate the mixture under shaking (150 rpm) for 1 hr at 37°C.
14. Spread 10 and 100 μl on LB agar plates containing the appropriate antibiotic and incubate at 37°C overnight.
15. Pick single colonies using toothpicks and cultivate to saturation in 5 ml LB medium containing antibiotic (37°C, 150 rpm) overnight.
16. Extract plasmid DNA using a QIAprep spin miniprep kit following the manufacturer's protocols. Elute the plasmid DNA in 50 μl elution buffer and quantify with a NanoDrop ND-1000 spectrophotometer. Check and confirm the fidelity of the construction by DNA sequencing (Eurofins).

Table 7 Amino Acid Composition of the Recombinant Protein Isoforms^a

Recombinant protein	Profiling amino acid sequence
TAT-HA- POI-EGFP-TEV-6×His	MDPKLG YGRKKRRQRRR GGSTMSGYPYDVPDYAGSMA[POI]VSKGEELFTGVVPILVELDGDVNGHKFSVSGEGEGDATYGKLT LKFI CTTGKLPVPWPTLVTTLT YGVQ CFSRYPDHMKQHDFFKSAMPEGYVQERTIFFKDDGNYKTRA EVKF EGDTLVNRIELKGIDFKEDGNILGHKLEYNYNSHNVYIMADKQKNGIKVNFKIRHNIEDGSVQLADHYQQNTPIGDGPVLLPDNH YLS TQSALSKDPNEKRDH MLLE FVTAAGITLGMD ELYKENLYFQG NSAVD HHHHHH
TAT-HA- POI-mCherry-TEV-6×His	MDPKLG YGRKKRRQRRR GGSTMSGYPYDVPDYAGSMA[POI]VSKGEEDNMA IIK EFMRFKVHMEGSVNGHEFEIEGEGEGRPYEGTQTAKLKVTKG GPLP FAWDILSPQFMYGSKAYVKHPADIPDYLKLSFPEGFKWERVMNFEDGGVVTVTQDSSLQDGEFIYKVKLRGTNFP SDGPVMQKKT MGWEASSERMY PEDG ALKGEIK QRLKL KDGGHYDAEVKTTYKAKKPVQLPGAYNVNIKLDITSHNEDYTIVEQY ERAEG RHSTGGMD ELYKENLYFQG NSAVD HHHHHH

^a For clarity the color code used for sequences is the same as that used in Figure 2 for fragments.

Transformation and selection procedure

17. Perform the transformation with the plasmid expressing the POI by using the heat shock method described in step 11. Thaw *E. coli* BL21(DE3) competent cells on ice.
18. Add 0.1 µg purified DNA plasmid to 50 µl competent cells and mix gently by pipetting up and down or flicking the tube 4 to 5 times (do not vortex).
19. After 20 min on ice, perform heat shock at 42°C for 45 s and immediately transfer cells on ice for 3 min.
20. Add 450 µl SOC medium and incubate the mixture under shaking (150 rpm) for 1 hr at 37°C.
21. Spread 10 and 100 µl of the transformation mixture on separate LB agar plates (Petri dishes) supplemented with the antibiotic.
22. Incubate overnight at 37°C.

Recombinant protein expression

23. Pick single colonies and cultivate to saturation in 5 ml LB containing antibiotic (37°C, 150 rpm) overnight.
24. Add pre-culture of transformed *E. coli* BL21(DE3) to 1 L of LB medium in 1-L glass Erlenmeyer flasks containing the antibiotic to reach a starting optical density at 600 nm (OD₆₀₀) of 0.1. Monitor the OD₆₀₀ until 0.6; then induce the recombinant POI expression (Table 7) for 4 h by adding 0.1% (w/v) L-arabinose (final concentration).
25. To ensure that the conditions for bacterial growth are optimal for the expression of the recombinant protein, monitor the bacterial growth by measuring the OD₆₀₀ during the culture and at the end of the expression phase (4 hr post-induction) (Fig. 4).
26. For cell harvesting, centrifuge the 1 L of LB culture medium 30 min at 6000 × g, 4°C, using the Avanti J-26S XP centrifuge (Fig. 5A).
27. Discard the supernatant LB and add an anti-protease complete EDTA-free protease inhibitor cocktail tablet to the bacterial pellet (one tablet per L culture medium).
28. Resuspend the bacterial pellet in 20 ml buffer A and store the 50-ml tube at –20°C overnight.




Figure 4 Growth profiles of empty *E. coli* BL21(DE3) and *E. coli* transformed with the pBAD-TAT-HA-POI-EGFP-TEV-6×His or pBAD-TAT-HA-POI-mCherry-TEV-6×His plasmids measured as function of time at OD₆₀₀ (induction of POI expression by L-arabinose addition is indicated by the arrow) (n = 3). POI, Protein of Interest.




Figure 5 Efficient cell lysis for protein extraction by sonication. **(A)** LB culture medium containing the *E. coli* BL21 cells after induction with L-arabinose (left) to promote protein production. Distribution of the culture medium in two centrifuge bottles before centrifugation (right) (Basic Protocol 1, step 26). **(B)** Optimizing protein extraction with the Bandelin Electronic Sonicator HD 4200 in cell lysis protocols (Basic Protocol 1, step 31). **(C)** Recovery of the bacterial pellet (right) and supernatant containing the POI (left) after centrifugation of bacteria following sonication (Basic Protocol 1, step 32). POI, Protein of Interest.

BASIC PROTOCOL 2

29. Thaw the frozen bacterial pellet by placing the 50-ml tube containing the bacterial pellet in a water bath at room temperature.
30. Add 100 μ l DNase I for every 1 L bacterial culture to degrade DNA that will be released by the bacterial lysis.
31. Perform bacterial lysis by sonication on ice, employing cycles of 30 s ON and 15 s OFF during 5 min, with the probe operating at 70% of maximal power (Fig. 5B).
32. Using a centrifuge and rotor 19776H, centrifuge the lysate 15 min at $12,000 \times g$, 4°C (Fig. 5C).
33. Filter the supernatant through a $0.45\text{-}\mu\text{m}$ filter. The supernatant containing the POI is ready for protein purification (Basic Protocol 2).

PROTEIN PURIFICATION

In this protocol, we describe the procedure of production and purification of recombinant protein isoforms (Fig. 1B). The remaining supernatant (from Basic Protocol 1, step 33) is then subjected to IMAC thanks to the $6\times$ His-tag. Subsequent wash steps remove impurities and then the POI is eluted using a buffer that disrupts the binding interaction. Further purification steps may include size-exclusion chromatography and/or ion-exchange chromatography to achieve higher purity. Throughout the protocol protein concentration is monitored and quality is assessed using SDS-PAGE and a brilliant Coomassie blue staining and by western blot. If conducted properly, the final purified POI should exhibit the expected molecular weight, high purity, and biological activity.

Materials

- Buffer A (see recipe)
- Buffer B (see recipe)
- Ultrapure H_2O
- Desalting buffer: Dulbecco's phosphate-buffered saline (DPBS), w/o Mg, w/o Ca (Dutscher, cat. no. L0615-500)
- Laemmli buffer, $5\times$ (see recipe)
- Polyacrylamide gels, 8% separating and 4% stacking (see recipes)
- SDS-PAGE buffer: 10% (v/v) Tris-glycine-SDS $10\times$ (Euromedex, cat. no. EU0510) in distilled H_2O
- PageRuler prestained protein ladder (Thermo Fisher Scientific, cat. no. SM0671)
- Quick Coomassie stain (NeoBiotech, cat. no. NB-45-00078-1litre)
- Transfer buffer: 10% (v/v) Tris-glycine $10\times$ (Euromedex, cat. no. EU0550) in distilled water, 20% (v/v) methanol (Carlo Erba, cat. no. 524102)
- Ponceau S red solution (Sigma-Aldrich, cat. no. P7170)
- TBS-Tween buffer (see recipe)
- Blocking buffer (see recipe)
- Mouse monoclonal antibodies:
 - Anti-HA-tag (F-7) (Santa Cruz Biotechnology, cat. no. sc-7392)
 - His-Tag (HIS. H8 / EH158) (Covalab, cat. no. mab90001-P)
 - Anti-POI
 - Anti-EGFP (Roche, cat. no. 45-11814460001)
- ECL anti-mouse IgG (from sheep), horseradish peroxidase-linked whole secondary antibodies (Fisher Scientific, cat. no. NA931V)
- SuperSignal West Pico PLUS (Thermo Scientific, cat. no.34577) or SuperSignal West Femto, maximum sensitivity substrate (Thermo Scientific, cat. no.34095).
- Preparative chromatography system (e.g., ÄKTA start protein purification system associated with UNICORN start control software)



Figure 6 Overview of the protein purification and desalting process. The protein sample containing the POI is injected using an automated injection system (indicated by the red arrowhead) and shown in the inset above-right (sample loading position). The bottles for buffer A (lysis buffer) and buffer B (elution buffer) are in the central polystyrene box. Fractions are collected using an automatic collector at the right of the picture. The process is monitored by measuring absorbance at a specific wavelength of 280 nm thanks to a computer at the left. The device is shown in its desalting configuration: the HiTrap Desalting is indicated by the blue arrowhead.

HisTrap fast flow, 1- or 5-ml, Cytiva (Dutscher, cat. no. 17-5255-01)
 HiTrap desalting, 5-ml, Cytiva (Dutscher, cat. no. 17-1408-01)
 Vivaspin 6, centrifugal concentrator, 50,000 MWCO PES (Grosseron, cat. no. VS0632)
 Benchtop centrifuge
 Heat block (Eppendorf ThermoMixer)
 Buffer tank lid
 Electrode assembly
 Combs electrophoresis Cell, 10-well, 1.5-mm
 Hamilton syringe, 50- μ l with needles
 Power supply generator
 Rotating platform
 Gel cassette
 Gel holder cassette and foam pads
 0.45- μ m pore nitrocellulose membrane (Amersham, cat. no. 10600016)
 Whatman paper sheets, 6 \times 9.5 cm, and sponges
 Electroblotting system
 Fusion Solo machine (Vilber Lourmat) for western blot detection

NOTE: It is possible that the POI aggregates and forms inclusion bodies that are difficult to solubilize and extract from the bacteria. In that case we recommend to process via mechanical rupture of the bacteria by sonication or by using a French press.

Protein fractionation

To purify your recombinant TAT-HA-POI-EGFP/mCherry-TEV-6 \times His protein by IMAC (exploiting the presence of a C-terminal 6 \times His-tag), use a simple benchtop nickel affinity column or combine with a preparative chromatography system, such as the ÄKTA Start system associated with a ÄKTA Start Fraction Collector “Frac30” and a UNICORN Start Software (Fig. 6). This can be carried out by employing an automated protein purification workflow with features like automated sample injection, fractions collection, and real-time monitoring caution.

1. Prepare bottles for buffer A, buffer B, ultrapure water, and the column HisTrap fast flow.
2. Set the IMAC purification program to use both buffer A and buffer B.
3. Equilibrate the column with 20 to 35 ml buffer A.

This step involves filling the column and allowing the buffer to flow through until it is stable and free of air bubbles.

4. Inject your protein sample containing the POI into the equilibrated column (usually done using an automated injection system).
5. Wash the column with 25 ml buffer A to remove unbound proteins and contaminants.
6. Start the imidazole concentration gradient between buffer A and buffer B, up to 100% buffer B, to ensure the controlled release of the POI from the column.
7. Collect 1 ml fractions from the column. Monitor the elution process in real-time by measuring absorbance at a specific wavelength of 280 nm (detection of proteins).

Protein desalting and concentration

8. Use a bottle of 1 × DPBS and a HiTrap desalting column.
9. Set the protein desalting program to only use pump A.
10. In the injection loop, re-inject the fraction containing your purified POI -1 ml per 1 ml and re-collect 5 to 10 desalted fractions of 1 ml each.
11. Concentrate the proteins (sample of 4 ml POI) using a Vivaspin 50 kDa cut-off through successive centrifugations of 5 min at $3500 \times g$, 4°C (Fig. 7A). Prior to use, rinse the concentrator column with water and subsequently with DPBS. Gently mix with a pipette before and after each centrifugation.

Take care not to exceed the specified speed indicated on the centrifugal filters.

12. Quantify protein concentration by measuring absorbance at 280 nm (extinction coefficients of 93,170 and 105,660 $M^{-1}cm^{-1}$ for the EGFP and the mCherry forms of the POI, respectively).

Analysis of recombinant protein expression by SDS-PAGE (Fig. 7B)

13. Load 10 to 20 µg of purified proteins (mixed and boiled 1/5 volume of 5 × Laemmli buffer) by using a Hamilton syringe on an 8% SDS-PAGE gel. Also load 5 µl molecular weight markers (typically ranging from 250 to 10 kDa).

The reticulation of the separating gel must be adjusted according to the molecular weight of the POI. Use a high reticulation for a low molecular weight protein (e.g., 12.5% reticulation for a 40 to 50 kDa protein) and a lower reticulation for a higher molecular weight protein (e.g., 8% reticulation for a 100 to 120 kDa protein).

14. Using a power supply generator, run at 30 mA in 1 × Tris-glycine-SDS buffer until the bromophenol blue line reaches the bottom of the glass plates (~2 hr).
15. For Coomassie blue staining (Figs. 7C and 7D), after migration carefully transfer the gel to a plastic or glass tray.
16. Cover the gel with Quick Coomassie stain with enough volume to allow it to move freely in the tray.
17. Cover with aluminum wrap and place on a platform rotator overnight at room temperature.




Figure 7 Protein concentration and characterization by SDS-PAGE. **(A)** Concentration of the POI by using a Vivaspin 6 type centrifugal concentrator. **(B)** Characterization of the recombinant POI expression by using an SDS-PAGE device. **(C and D)** Assessed of POI production and purity by Coomassie blue staining and western blotting. Sample A corresponds to the TAT-HA-POI-EGFP-TEV-6×His protein and sample B to the TAT-HA-POI-mCherry-TEV-6×His protein. Molecular weight markers are indicated on the left part of each panel (in kDa).

18. Proceed to the gel destaining step by multiple washes with ultrapure water until the band corresponding to the POI is easily visible.
19. For western blotting (Figs. 7C and 7D), transfer the gel onto nitrocellulose and proceed to electro-blotting in a device dedicated for that purpose (e.g., BioRad blotting system).
20. After transfer, check the transfer efficiency (and if necessary, the equal loading of the samples) by incubating the nitrocellulose membrane in the Ponceau S red solution for 1 min.
21. Discard the excess of solution and wash the membrane with ultrapure water until the band(s) is(are) easily visible.
22. Take a picture of the stained membrane for archiving.
23. Destain the membrane by incubation in the TBS-Tween buffer for a few seconds until gentle shaking.
24. Saturate the membrane by incubation in the blocking buffer for 45 min at room temperature (use a rotating platform for gentle shaking).
25. Discard the blocking buffer and incubate the membrane with the appropriate antibodies (mouse polyclonal anti-HA, mouse monoclonal anti-6×His, mouse monoclonal anti-POI at the appropriate dilution according to the manufacturer's instructions in blocking buffer).
26. Place the membrane on a rotating platform (gentle shaking) overnight at 4°C.

27. Wash the membranes with TBS-Tween 3 times for 7 min each and then incubate with an anti-mouse horseradish peroxidase-labeled secondary antibody (usually at a dilution of 1/10,000) for 1 hr.
28. Wash the membranes with TBS-Tween 3 times for 7 min each.
29. Add a mixed volume of buffers A and B (1:1) of SuperSignal West Pico Plus onto a parafilm sheet placed on the bench.
30. Position the nitrocellulose membrane so that the proteins directly face the revelation liquid. Wait for 5 min in the dark.
31. Transfer the membrane onto a tray for protein detection and place the tray into the Fusion Solo machine.
32. For capturing images using enhanced chemiluminescence, use the Fusion software (chemiluminescence and fluorescence imaging).

In case your POI expression is low, or the signal is weak with the SuperSignal West Pico plus, we recommend using the SuperSignal West Femto, Maximum sensitivity substrate, and repeat the same procedure.

Protein storage

33. According to protein stability, store the POI at -80°C in 10% (v/v) sterile glycerol until use.

BASIC PROTOCOL 3

PROTEIN TRANSDUCTION IN MAMMALIAN CELLS

In this protocol, we describe the procedure of protein transduction within Hep3B mammalian cells. The investigation of protein entry into mammalian cells is a pivotal aspect of cellular and molecular biology, shedding light on essential physiological processes and enabling the development of targeted therapeutic strategies. This protocol delineates a fundamental approach for the transduction of proteins into mammalian cells, a process to efficiently deliver exogenous recombinant proteins. Furthermore, this protocol provides a detailed guide for monitoring the intracellular fate of transduced proteins through immunofluorescence microscopy, offering a powerful tool for visualizing and analyzing their subcellular localization and dynamics through fluorescent tags, such as EGFP and mCherry.

Materials

- Hep3B cells (ATCC, cat. no. HB-8064)
- Gibco minimum essential medium (MEM) (Dutscher, cat. no. L0430-50), or other cell culture medium adapted to the cell line used
- Fetal calf serum (FCS) (Corning, cat. no. 35-079-CV)
- 2 mM L-glutamine (Gibco by Life Technologies, cat. no. 25030-024)
- 1 mM sodium pyruvate (Gibco by Life Technologies, cat. no. 11360070)
- 50 mg/L antibiotics (streptomycin, penicillin) (Gibco by Life Technologies, cat. no. 15070-063)
- DPBS w/o Mg, w/o Ca (Dutsher, cat. no. L0615-500)
- Trypsin-EDTA, 1× in PBS w/o Ca, w/o Mg w/o phenol red (Dutsher, cat. no. L0940-100)
- Trypan blue solution (Sigma-Aldrich, cat. no. T8154)
- Purified POI in DPBS, 1×
- DAPI (4',6-diamidino-2-phenylindole dihydrochloride) (Sigma-Aldrich, cat. no. D9542)
- 70% ethanol
- Moviol mounting medium (see recipe)
- Water bath
- T25 flasks (Corning, cat. no. 430720U)

Humidified cell culture incubator enriched with 5% (v/v) CO₂
 Leica DM IRB optical microscope
 T75 flasks (Corning, cat. no. 430641U)
 Suction pump
 15-ml sterile conical tubes (Falcon, cat. no. 352097)
 Malassez slide
 Cell counter
 24-well plates (Corning, cat. no. 3526)
 120-mm coverslips
 4% (w/v) paraformaldehyde (PAF) (Sigma-Aldrich, cat. no. P6148)
 Humid chamber
 Whatman paper
 Parafilm
 Zeiss LSM 780 confocal Microscope
 Fiji ImageJ

CAUTION: Work consistently under the hood (with ventilation) and always wear gloves when handling PAF due to its toxic properties as a CMR substance.

NOTE: In cell culture, it is essential to maintain a sterile working environment. Clean the biological safety cabinet (BSC) thoroughly with 70% ethanol and disinfectant solution before each manipulation or experiment. Thoroughly clean all bottles used in cell culture with 70% ethanol before handling, and always manipulate under the BSC to prevent potential contamination.

Cell culture

1. Prepare the cell culture medium by mixing MEM to 10% (v/v) FCS, 2 mM L-glutamine, 1 mM sodium pyruvate, and 50 mg/L antibiotics. Place the cell culture medium, 1 × DPBS, and trypsin in a water bath at 37°C for ~10 to 20 min to achieve the optimal temperature for cell culture maintenance.
2. Initiate the culture of Hep3B cells in a T25 flask within an incubator set at 37°C with 5% (v/v) CO₂.
3. Monitor cell proliferation using an optical (visible light) microscope to ensure viability; prevent contamination before the experiment and maintain confluency while adjusting the medium if necessary.
4. Once Hep3B cells reach 70% confluence in a T75 flask, discard the cell medium.
5. Wash the cells with DPBS and discard DPBS using a suction pump.
6. Add 2 ml trypsin and place the flask in the cell incubator at 37°C until the cells detach from the flask (~5 to 10 min). Confirm cell detachment under the optical microscope.
7. Add 4 ml cell culture medium, twice the volume of trypsin on the cell monolayer. Discard the total 6 ml and transfer it to a sterile 15-ml sterile conical tube.
8. Take a 10 µl cell sample and mix with 10 µl trypan blue (1:1 ratio).
9. Use a Malassez slide for cell counting, placing 10 µl in each chamber and scan with a cell counting machine. Calculate the volume of cells to add to the culture medium for a final volume of 1 ml per well.

Protein transduction within living cells

10. In a 24-well plate, place a 120-mm coverslip in each well. Seed the cells for 24 hr by depositing ~37,500 cells with cell culture medium to achieve a final volume of 1 ml per well.

Anny et al.

17 of 26

11. After 24 hr, the cells should be well adherent and have reached $\sim 70\%$ confluence.
12. Discard 1 ml medium, then add 1 to 5 μM of purified TAT-HA-POI-XFP (or POI-XFP if the TEV sequence has been cleaved) onto cells with the required culture medium volume to reach a final volume of 1 ml per well. Incubate the cells at 37°C for 2 hr.
13. Discard the medium, perform three quick washes with cold $1\times$ DPBS, and fix the cells on the coverslips by adding 500 μl of 4% (w/v) PAF per well for 30 min at room temperature.
We can consider that cell transduction has occurred after 2 hr, otherwise adjust the time of contact between the POI, which the concentration can also be modulated, and the cells.
Do not exceed a 30 min exposure to 4% (w/v) PAF, as it is toxic to cells.
14. Wash the cells three times with DBPS for 5 min each.
Conduct the initial cell wash with $1\times$ DPBS under the hood and in ventilation mode to eliminate any residual 4% (w/v) PAF.
15. Meanwhile, to avoid a drying out of the coverslips, prepare a humid chamber using a support, place a Whatman paper on it, wet it with ultrapure water, and cover it with parafilm.
16. Place the coverslip in the humid chamber and add 100 μl of $1\times$ DPBS to cover the cells and prevent drying.
17. Discard DBPS and add DAPI to stain nuclei directly to each coverslip for 10 min at room temperature in three washes of 5 min each. Clean the coverslip with 70% ethanol and let it dry.
18. Add 5 μl Moviol (mounting liquid) to the coverslip and place it directly on the Moviol so that the side containing the fixed cells faces the liquid.
19. Let the coverslip dry at room temperature overnight in the dark. The next day, the coverslip is ready for observation under a confocal microscope.
20. Capture the images by using a confocal microscope:
 - a. Channel 1 (DAPI), λ_{ex} 405 nm, λ_{em} 410 to 470 nm.
 - b. Channel 2 (EGFP), λ_{ex} 488 nm, λ_{em} 500 to 543 nm.
 - c. Channel 3 (mCherry), λ_{ex} 561 nm, λ_{em} 570-615 nm.
21. Images can be analyzed by using Fiji ImageJ software (Fig. 8).

REAGENTS AND SOLUTIONS

Blocking buffer

5% (w/v) non-fatty-acid milk (for food use) or BSA (bovine serum albumin) (Sigma-Aldrich, cat. no. A9647) in TBS-Tween buffer (see recipe)
 Mix well
 Prepare fresh

Buffer A

3.15 g Tris (20 mM final; Euromedex, cat. no. EU001B)
 17.53 g NaCl (300 mM final; Carlo Erba, cat. no. 479685)
 1.36 g imidazole (20 mM final; Sigma-Aldrich, cat. no. I202)
 Adjust to 1 L with ultrapure H_2O
 Adjust to pH 8.0 with HCl (37% w/v) (Carlo Erba, cat. no. 403871)
 Mix well and filter using a 0.22- μm filter
 Store up to 1 month at 4°C




Figure 8 Fluorescence microscopy after treatment for 2 hr of Hep3B cells with 2 μ M of recombinant POI-EGFP and POI-mCherry fusion isoforms. Cells were fixed on coverslips with 4% (w/v) paraformaldehyde. Cells were counterstained for DNA with 1 μ g/ml DAPI (blue fluorescence), washed with PBS and then mounted in Mowiol for microscopy analysis. Scale bar: 20 μ m.

Buffer B

3.15 g Tris (20 mM final; Euromedex, cat. no. EU001B)
 17.53 g NaCl (300 mM final; Carlo Erba, cat. no. 479685)
 34 g imidazole (500 mM final; Sigma-Aldrich, cat. no. I202)
 Adjust to 1 L with ultrapure H₂O
 Adjust to pH 8.0 with HCl (37% w/v) (Carlo Erba, cat. no. 403871)
 Mix well and filter using a 0.22- μ m filter
 Store up to 1 month at 4°C

Anny et al.

19 of 26

Laemmli buffer, 5×

12.11 g Tris (250 mM final; Euromedex, cat. no. EU001B)
 5% (w/v) SDS (Sigma-Aldrich, cat. no. L3771)
 5% (v/v) 2-mercaptoethanol (Sigma-Aldrich, cat. no. M3148)
 40% (v/v) glycerol (Sigma-Aldrich, cat. no. G9012)
 0.02% (w/v) bromophenol blue (Sigma-Aldrich, cat. no. B0126)
 Bring up to 1 L with ultrapure H₂O
 Adjust to pH 8.0 with HCl (37% w/v) (Carlo Erba, cat. no. 403871)
 Mix well
 Store at up to 1 month at 4°C

Polyacrylamide separating gel, 8%

3.75 ml of 1.5 M Tris·HCl pH 8.8 (Current Protocols, 2006)
 50 µl of 20% (w/v) SDS (Sigma-Aldrich, cat. no. L3771)
 2 ml of 40% (w/v) acrylamide/bis-acrylamide (ratio 29:1) (Sigma-Aldrich, cat. no. A2792)
 30 µl Temed (*N,N,N',N'*-tetramethylethylenediamine) (Sigma-Aldrich, cat. no. T9281)
 30 µl of 10% (w/v) ammonium persulfate (Sigma-Aldrich, cat. no. A7460)
 Bring up to 10 ml with ultrapure H₂O
 Mix well
 Prepare fresh

Polyacrylamide stacking gel, 4%

1.5 ml of 0.5 M Tris·HCl pH 6.8 (Current Protocols, 2006)
 50 µl of 20% (w/v) SDS (Sigma-Aldrich, cat. no. L3771)
 1 ml of 40% (w/v) acrylamide/bis-acrylamide (ratio 29:1) (Sigma-Aldrich, cat. no. A2792)
 60 µl Temed (*N,N,N',N'*-tetramethylethylenediamine) (Sigma-Aldrich, cat. no. T9281)
 50 µl of 10% (w/v) ammonium persulfate (Sigma-Aldrich, cat. no. A7460)
 Bring up to 10 ml with ultrapure H₂O
 Mix well
 Prepare fresh

Mowiol mounting medium

6 g glycerol
 2.4 g mowiol 4-88 (biovalley, cat. no. 17951-500)
 6 ml distilled H₂O
 12 ml 0.2 M Tris·HCl pH 8.5 (Current Protocols, 2006)
 Mix well and centrifuge 10 min at 1500 × g, room temperature, to clarify the solution
 Store up to 1 month at 4°C or 1 year at −20°C

TBS-Tween buffer

4.2 g Tris (15 mM final; Euromedex, cat. no. EU001B)
 8.1 g NaCl (300 mM final; Carlo Erba, cat. no. 479685)
 0.05% (v/v) Tween 20 (Fisher Scientific, cat. no. M12247)
 Adjust to 1 L with distilled H₂O
 Adjust to pH 8.0 with HCl (37% w/v) (Carlo Erba, cat. no. 403871)
 Mix well
 Store up to 1 week at room temperature

COMMENTARY

Critical Parameters

Several steps in this experimental strategy are critical and therefore can greatly influence its success. It is important to correctly design the primers so that the different PCRs that precede the DNA molecular assembly are as specific as possible to generate a single band visualized on the agarose gel (Basic Protocol 1). This step is fundamental to getting off to a good start. Another crucial point is the bacteria subculturing stage. It is important to pick well-isolated bacterial colonies back into liquid culture so as to not mix different clones (Basic Protocol 1). The last step concerning the transduction of cells is relatively simple because the use of penetration peptides makes it possible to avoid many of the steps inherent in the transfection of plasmid vectors (Basic Protocol 3). Nevertheless, it is important here to test variable quantities of proteins to be transduced to optimize their subsequent study. If the POI is not used quickly after its purification, it is extremely important to store it in suitable conditions. Nevertheless, we encourage users to extemporaneously prepare the protein to transduce it immediately after purification (see Table 8). Other potential critical parameters are listed in Table 8.

Troubleshooting Table

Table 8 lists the most common problems that may be encountered during protein production detailed in this article (Table 8).

Statistical Analysis

GraphPad Prism 9 or any other statistics tool can be used for calculating the standard deviations of the growth curves of bacteria, as those shown in Figure 4.

Understanding Results

Basic Protocols 1 and 2

The data obtained by Basic Protocols 1 and 2 generate agarose gel electrophoretic profiles to assess the quantity and quality of the PCR products, and polyacrylamide gel profiles that can be visualized either by treatment of the gels with a solution of Coomassie blue or by electro-blotting of the resolved proteins by SDS-PAGE onto nitrocellulose. These experiments require carefully supporting the samples to be analyzed with size markers for agarose gels and molecular mass markers for polyacrylamide gels; be careful not to confuse the types of markers, the first being expressed in base pairs (bp), the second in kilodaltons (kDa).

Table 8 Troubleshooting Table for Protein Production

Problem	Possible Cause	Solution
Poor protein stability	Inappropriate storage conditions or buffer	Store protein at recommended temperature; optimize buffer composition; add proteases inhibitors to buffers
Low protein yield	Inefficient cell lysis or purification method	Optimize lysis conditions; consider alternative purification methods
Contaminants in elution	Carryover from previous steps or column issues	Ensure proper column equilibration; optimize washing steps
Low/no signal in western blot results	Poor antibody binding	Ensure primary and secondary antibodies are compatible and correctly diluted
High background or low ratio expected signal/background in western blot	Antibodies too much interact with the membrane or with the blocking reagent	Increase the time and the number of washings; change the blocking reagent; increase the dilution of the primary or/and the secondary antibodies
Weak or no fluorescence signal	Too low protein concentration in contact with the cell	Increase the concentration of the protein
Cell autofluorescence	Presence of residual medium or contaminants	Ensure thorough washing after treatment
Protein aggregation	Improper storage conditions or handling	Store protein appropriately; handle with care

The presentation of the data requires labeling the figures by clearly identifying at the top of the gels the samples loaded on the left or the right lateral side, and the values corresponding to each mark of size (bp) or molecular mass (kDa). By comparing the migration distance of the band corresponding to the PCR product or to the protein produced, it is possible to prove not only that the product is indeed present, but also to estimate its abundance and purity. The western blot provides an advantage since the use of antibodies guarantees an additional argument simply because the apparent molecular mass is indeed what is expected. Make sure to check the supplier's information sheet whether these antibodies can be used for western blotting. Some antibodies are more suitable for immunoprecipitation, ELISA, or immunofluorescence since they recognize proteins in their native form while others are better for western blot because they were generated against a short peptide sequence. This is the case for the latter in this series of protocols, we used antibodies directed against tag "labels" (HA and 6×His), which generally guarantees their use in western blotting, while anti-HA antibodies are better for western blotting than anti-6×His antibodies preferentially used for protein purification. The problem of non-recognition can, however, be encountered with anti-POI antibodies. In fact, in Figure 3 and Figure 7C-D, we can consider on the one hand the good purity of the different PCR products (this must be ensured before moving on to the molecular cloning stage, which will be more perilous if the PCRs are not sufficiently pure) and the protein thus produced. It is common to observe, as we do in Figure 7C-D, bands of molecular mass lower than the expected band; these are generally proteolytic degradation bands generated during production and purification. In our case it turns out that the mCherry form of our POI is less stable than the GFP form (it may be useful in this case to use larger quantities of protease inhibitors during the process).

Basic Protocol 3

Regarding images obtained by microscopy in confocal mode (Basic Protocol 3), they must be clearly labeled by indicating the corresponding condition. It is very useful to merge with the DAPI image, which allows to locate the POI at a glance. Our microscopy images show a cytoplasmic distribution of our POI in both the EGFP and mCherry versions. Finally, a scale bar (20 μm in our experience) makes it possible to judge the distances on

the microscopy images. It is common and even advisable to show separate images and merge for immunofluorescence microscopy experiments.

Time Considerations

Total time required to complete the full experimental strategy: 3 to 4 weeks.

Basic Protocol 1 (7 days)

Amplification of the various PCR products

Samples preparation: 10 min

PCR: 3 hr

Agarose gel preparation and polymerization: 30 min

Agarose gel electrophoresis: 30 min

Ethidium bromide staining: 15 min

Bands visualization by using UV transilluminator: 10 min

Bands excision by using UV transilluminator: 15 min

PCR gel purification: 1 hr

Total: 1 day

DNA molecular assembly

Plasmid assembly procedure: 1 hr

Transformation and selection procedure (*E. coli* 5-alpha): 1 days

DNA extraction and purification: 1 hr

DNA sequencing: 2 days

Total: 4 days

Transformation and selection procedure in E. coli BL21(DE3)

Total: 1 day

Recombinant protein expression

Total: 1 day

Basic Protocol 2 (5 days)

Protein fractionation

Total: 1 day

Protein desalting, concentration, and quantification

Total: 3 days

Analysis of recombinant protein expression by SDS-PAGE

Coomassie blue staining

Western blot

Total: 1 day

Basic Protocol 3 (3 days)

Cell seeding

Total: 1 day

Protein transduction
Protein transduction: 2 hr
Slide and coverslip preparation
Total: 1 day

Data collection and analysis
Immunofluorescence microscopy: data
collecting
Data analysis: 1 to 2 hr
Total: 1 day

Acknowledgments

The authors are grateful to the European Union's Horizon 2020 research and innovation program that funded this work (Marie Skłodowska-Curie grant agreement No 847568), the Centre National de la Recherche Scientifique (CNRS) and the University of Lille for their recurrent fundings, and the Plateformes Lilloises en Biologie et Santé (PLBS) for help in using fluorescent microscopy.

Author Contributions

Christer Abou Anny: Conceptualization; data curation; formal analysis; methodology; writing original draft. **Sébastien Nouaille:** Conceptualization; investigation; methodology; supervision; validation; writing review and editing. **Régis Fauré:** Conceptualization; investigation; methodology; supervision; validation; writing review and editing. **Céline Schulz:** Conceptualization; formal analysis; methodology; writing review and editing. **Corentin Spriet:** Conceptualization; methodology; writing review and editing. **Isabelle Huvent:** Conceptualization; investigation; methodology; supervision; writing review and editing. **Christophe Biot:** Conceptualization; formal analysis; funding acquisition; investigation; resources; supervision; validation; writing review and editing. **Tony Lefebvre:** Conceptualization; formal analysis; funding acquisition; investigation; project administration; resources; supervision; writing original draft.

Conflict of Interest

The authors declare no conflict of interest.

Data Availability Statement

Data sharing is not applicable to this article as no new data were created in this study.

Literature Cited

Ai, H., Baird, M. A., Shen, Y., Davidson, M. W., & Campbell, R. E. (2014). Engineering and characterizing monomeric fluorescent proteins for live-cell imaging applications. *Nature Pro-*

ocols, 9(4), 910–928. <https://doi.org/10.1038/nprot.2014.054>

Andersen, D. C., & Krummen, L. (2002). Recombinant protein expression for therapeutic applications. *Current Opinion in Biotechnology*, 13(2), 117–123. [https://doi.org/10.1016/S0958-1669\(02\)00300-2](https://doi.org/10.1016/S0958-1669(02)00300-2)

MacKay, A. J., & Szoka, F. C. Jr. (2003). HIV TAT protein transduction domain mediated cell binding and intracellular delivery of nanoparticles. *Journal of Dispersion Science and Technology*, 24(3–4), 465–473. <https://doi.org/10.1081/DIS-120021802>

Anné, J., Maldonado, B., Van Impe, J., Van Mellaert, L., & Bernaerts, K. (2012). Recombinant protein production and streptomycetes. *Journal of Biotechnology*, 158(4), 159–167. <https://doi.org/10.1016/j.jbiotec.2011.06.028>

Assenberg, R., Wan, P. T., Geisse, S., & Mayr, L. M. (2013). Advances in recombinant protein expression for use in pharmaceutical research. *Current Opinion in Structural Biology*, 23(3), 393–402. <https://doi.org/10.1016/j.sbi.2013.03.008>

Baneyx, F. (1999). Recombinant protein expression in *Escherichia coli*. *Current Opinion in Biotechnology*, 10(5), 411–421. [https://doi.org/10.1016/S0958-1669\(99\)00003-8](https://doi.org/10.1016/S0958-1669(99)00003-8)

Baneyx, F., & Mujacic, M. (2004). Recombinant protein folding and misfolding in *Escherichia coli*. *Nature Biotechnology*, 22(11), 1399–1408. <https://doi.org/10.1038/nbt1029>

Barnes, L. M., Bentley, C. M., & Dickson, A. J. (2003). Stability of protein production from recombinant mammalian cells. *Biotechnology and Bioengineering*, 81(6), 631–639. <https://doi.org/10.1002/bit.10517>

Becker-Hapak, M., McAllister, S. S., & Dowdy, S. F. (2001). TAT-mediated protein transduction into mammalian cells. *Methods*, 24(3), 247–256. <https://doi.org/10.1006/meth.2001.1186>

Block, H., Maertens, B., Priestersbach, A., Brinker, N., Kubicek, J., Fabis, R., Labahn, J., & Schäfer, F. (2009). Immobilized-metal affinity chromatography (IMAC): A review. *Methods in Enzymology*, 463, 439–473. [https://doi.org/10.1016/S0076-6879\(09\)63027-5](https://doi.org/10.1016/S0076-6879(09)63027-5)

Bordat, A., Houvenaghel, M.-C., & German-Retana, S. (2015). Gibson assembly: An easy way to clone potyviral full-length infectious cDNA clones expressing an ectopic VPg. *Virology Journal*, 12(1), 89. <https://doi.org/10.1186/s12985-015-0315-3>

Brondyk, W. H. (2009). Selecting an appropriate method for expressing a recombinant protein. *Methods in Enzymology*, 463, 131–147. [https://doi.org/10.1016/S0076-6879\(09\)63011-1](https://doi.org/10.1016/S0076-6879(09)63011-1)

Burnett, M. J. B., & Burnett, A. C. (2020). Therapeutic recombinant protein production in plants: Challenges and opportunities. *Plants, People, Planet*, 2(2), 121–132. <https://doi.org/10.1002/ppp3.10073>

Casini, A., Storch, M., Baldwin, G. S., & Ellis, T. (2015). Bricks and blueprints: Methods and

- standards for DNA assembly. *Nature Reviews Molecular Cell Biology*, 16(9), 568–576. <https://doi.org/10.1038/nrm4014>
- Chen, R. (2012). Bacterial expression systems for recombinant protein production: *E. coli* and beyond. *Biotechnology Advances*, 30(5), 1102–1107. <https://doi.org/10.1016/j.biotechadv.2011.09.013>
- Cohen, S. N. (2013). DNA cloning: A personal view after 40 years. *Proceedings of the National Academy of Sciences*, 110(39), 15521–15529. <https://doi.org/10.1073/pnas.1313397110>
- Cregg, J. M., Cereghino, J. L., Shi, J., & Higgins, D. R. (2000). Recombinant protein expression in *Pichia pastoris*. *Molecular Biotechnology*, 16(1), 23–52. <https://doi.org/10.1385/MB:16:1:23>
- Current Protocols. (2006). Commonly Used Reagents. *Current Protocols in Microbiology*, 00, A.2A.1–A.2A.15. <https://doi.org/10.1002/9780471729259.mca02as00>
- Eguchi, A., Akuta, T., Okuyama, H., Senda, T., Yokoi, H., Inokuchi, H., Fujita, S., Hayakawa, T., Takeda, K., Hasegawa, M., & Nakanishi, M. (2001). Protein transduction domain of HIV-1 Tat protein promotes efficient delivery of DNA into mammalian cells. *Journal of Biological Chemistry*, 276(28), 26204–26210. <https://doi.org/10.1074/jbc.M010625200>
- Farzaneh, M., Hassani, S.-N., Mozdziak, P., & Baharvand, H. (2017). Avian embryos and related cell lines: A convenient platform for recombinant proteins and vaccine production. *Biotechnology Journal*, 12(5), 1600598. <https://doi.org/10.1002/biot.201600598>
- Ferro, M. M. M., Ramos-Sobrinho, R., Xavier, C. A. D., Zerbini, F. M., Lima, G. S. A., Nagata, T., & Assunção, I. P. (2019). New approach for the construction of infectious clones of a circular DNA plant virus using Gibson assembly. *Journal of Virological Methods*, 263, 20–23. <https://doi.org/10.1016/j.jviromet.2018.10.017>
- Fittipaldi, A., & Giacca, M. (2005). Transcellular protein transduction using the Tat protein of HIV-1. *Advanced Drug Delivery Reviews*, 57(4), 597–608. <https://doi.org/10.1016/j.addr.2004.10.011>
- Ford, K. G., Souberbielle, B. E., Darling, D., & Farzaneh, F. (2001). Protein transduction: An alternative to genetic intervention? *Gene Therapy*, 8(1), 1–4. <https://doi.org/10.1038/sj.gt.3301383>
- Froger, A., & Hall, J. E. (2007). Transformation of plasmid DNA into *E. coli* using the heat shock method. *Journal of Visualized Experiments*, 6, e253. <https://doi.org/10.3791/253>
- Gaberc-Porekar, V., & Menart, V. (2001). Perspectives of immobilized-metal affinity chromatography. *Journal of Biochemical and Biophysical Methods*, 49(1), 335–360. [https://doi.org/10.1016/S0165-022X\(01\)00207-X](https://doi.org/10.1016/S0165-022X(01)00207-X)
- Gibson, D. G., Young, L., Chuang, R.-Y., Venter, J. C., Hutchison, C. A., & Smith, H. O. (2009). Enzymatic assembly of DNA molecules up to several hundred kilobases. *Nature Methods*, 6(5), 343–345. <https://doi.org/10.1038/nmeth.1318>
- Gopal, G. J., & Kumar, A. (2013). Strategies for the production of recombinant protein in *Escherichia coli*. *The Protein Journal*, 32(6), 419–425. <https://doi.org/10.1007/s10930-013-9502-5>
- Greenblatt, J., & Schleif, R. (1971). Arabinose C protein: Regulation of the arabinose operon in vitro. *Nature New Biology*, 233(40), 166–170. <https://doi.org/10.1038/newbio233166a0>
- Gump, J. M., & Dowdy, S. F. (2007). TAT transduction: The molecular mechanism and therapeutic prospects. *Trends in Molecular Medicine*, 13(10), 443–448. <https://doi.org/10.1016/j.molmed.2007.08.002>
- Gump, J. M., June, R. K., & Dowdy, S. F. (2010). Revised role of glycosaminoglycans in TAT protein transduction domain-mediated cellular transduction. *Journal of Biological Chemistry*, 285(2), 1500–1507. <https://doi.org/10.1074/jbc.M109.021964>
- Guo, Q., Zhao, G., Hao, F., & Guan, Y. (2012). Effects of the TAT peptide orientation and relative location on the protein transduction efficiency. *Chemical Biology & Drug Design*, 79(5), 683–690. <https://doi.org/10.1111/j.1747-0285.2011.01315.x>
- Helmuth, J. A., Burckhardt, C. J., Greber, U. F., & Sbalzarini, I. F. (2009). Shape reconstruction of subcellular structures from live cell fluorescence microscopy images. *Journal of Structural Biology*, 167(1), 1–10. <https://doi.org/10.1016/j.jsb.2009.03.017>
- Hopkins, R. F., Wall, V. E., & Esposito, D. (2012). Optimizing transient recombinant protein expression in mammalian cells. *Methods in Molecular Biology*, 801, 251–268. https://doi.org/10.1007/978-1-61779-352-3_16
- Jäger, V., Büssow, K., & Schirrmann, T. (2015). Transient recombinant protein expression in mammalian cells. In M. Al-Rubeai (Ed.), *Animal Cell Culture. Cell Engineering* (Vol. 9, pp. 27–64). Springer. https://doi.org/10.1007/978-3-319-10320-4_2
- Jakobs, S., Subramaniam, V., Schönle, A., Jovin, T. M., & Hell, S. W. (2000). EGFP and DsRed expressing cultures of *Escherichia coli* imaged by confocal, two-photon and fluorescence lifetime microscopy. *FEBS Letters*, 479(3), 131–135. [https://doi.org/10.1016/S0014-5793\(00\)01896-2](https://doi.org/10.1016/S0014-5793(00)01896-2)
- Jang, C.-W., & Magnuson, T. (2013). A novel selection marker for efficient DNA cloning and recombineering in *E. coli*. *PLoS ONE*, 8(2), e57075. <https://doi.org/10.1371/journal.pone.0057075>
- Joung, Y. H., Park, S. H., Moon, K.-B., Jeon, J.-H., Cho, H.-S., & Kim, H.-S. (2016). The last ten years of advancements in plant-derived recombinant vaccines against Hepatitis B. *International Journal of Molecular Sciences*, 17(10), 1715. <https://doi.org/10.3390/ijms17101715>

- Kabouridis, P. S. (2003). Biological applications of protein transduction technology. *Trends in Biotechnology*, 21(11), 498–503. <https://doi.org/10.1016/j.tibtech.2003.09.008>
- Kaplan, I. M., Wadia, J. S., & Dowdy, S. F. (2005). Cationic TAT peptide transduction domain enters cells by macropinocytosis. *Journal of Controlled Release*, 102(1), 247–253. <https://doi.org/10.1016/j.jconrel.2004.10.018>
- Kirchhausen, T. (2009). Imaging endocytic clathrin structures in living cells. *Trends in Cell Biology*, 19(11), 596–605. <https://doi.org/10.1016/j.tcb.2009.09.002>
- Kostylev, M., Otwell, A. E., Richardson, R. E., & Suzuki, Y. (2015). Cloning should be simple: *Escherichia coli* DH5 α -mediated assembly of multiple DNA fragments with short end homologies. *PLoS ONE*, 10(9), e0137466. <https://doi.org/10.1371/journal.pone.0137466>
- Leifert, J. A., Harkins, S., & Whitton, J. L. (2002). Full-length proteins attached to the HIV tat protein transduction domain are neither transduced between cells, nor exhibit enhanced immunogenicity. *Gene Therapy*, 9(21), 1422–1428. <https://doi.org/10.1038/sj.gt.3301819>
- Li, C., Ji, C., Huguet-Tapia, J. C., White, F. F., Dong, H., & Yang, B. (2019). An efficient method to clone TAL effector genes from *Xanthomonas oryzae* using Gibson assembly. *Molecular Plant Pathology*, 20(10), 1453–1462. <https://doi.org/10.1111/mpp.12820>
- Li, L., Jiang, W., & Lu, Y. (2018). A modified Gibson assembly method for cloning large DNA fragments with high GC contents. *Methods in Molecular Biology*, 1671, 203–209. https://doi.org/10.1007/978-1-4939-7295-1_13
- Li, Z.-N., Jelkmann, W., Sun, P., & Zhang, L. (2020). Construction of full-length infectious cDNA clones of apple stem grooving virus using Gibson assembly method. *Virus Research*, 276, 197790. <https://doi.org/10.1016/j.virusres.2019.197790>
- Mattanovich, D., Branduardi, P., Dato, L., Gasser, B., Sauer, M., & Porro, D. (2012). Recombinant protein production in yeasts. *Methods in Molecular Biology*, 824, 329–358. https://doi.org/10.1007/978-1-61779-433-9_17
- McDonald, D., Vodicka, M. A., Lucero, G., Svitkina, T. M., Borisy, G. G., Emerman, M., & Hope, T. J. (2002). Visualization of the intracellular behavior of HIV in living cells. *Journal of Cell Biology*, 159(3), 441–452. <https://doi.org/10.1083/jcb.200203150>
- McRae, S. R., Brown, C. L., & Bushell, G. R. (2005). Rapid purification of EGFP, EYFP, and ECFP with high yield and purity. *Protein Expression and Purification*, 41(1), 121–127. <https://doi.org/10.1016/j.pep.2004.12.030>
- Newman, R. H., Fosbrink, M. D., & Zhang, J. (2011). Genetically encodable fluorescent biosensors for tracking signaling dynamics in living cells. *Chemical Reviews*, 111(5), 3614–3666. <https://doi.org/10.1021/cr100002u>
- Nienhaus, K., & Nienhaus, G. U. (2014). Fluorescent proteins for live-cell imaging with super-resolution. *Chemical Society Reviews*, 43(4), 1088–1106. <https://doi.org/10.1039/C3CS60171D>
- Noguchi, H., & Matsumoto, S. (2006). Protein transduction technology: A novel therapeutic perspective. *Acta Medica Okayama*, 60(1), 1–11. <http://doi.org/10.18926/AMO/30757>
- Nolte, C., Matyash, M., Pivneva, T., Schipke, C. G., Ohlemeyer, C., Hanisch, U.-K., Kirchhoff, F., & Kettenmann, H. (2001). GFAP promoter-controlled EGFP-expressing transgenic mice: A tool to visualize astrocytes and astrogliosis in living brain tissue. *Glia*, 33(1), 72–86.
- Norris, A. D., Kim, H.-M., Colaiácovo, M. P., & Calarco, J. A. (2015). Efficient genome editing in *Caenorhabditis elegans* with a toolkit of dual-marker selection cassettes. *Genetics*, 201(2), 449–458. <https://doi.org/10.1534/genetics.115.180679>
- Overton, T. W. (2014). Recombinant protein production in bacterial hosts. *Drug Discovery Today*, 19(5), 590–601. <https://doi.org/10.1016/j.drudis.2013.11.008>
- Pawson, T., & Nash, P. (2000). Protein-protein interactions define specificity in signal transduction. *Genes & Development*, 14(9), 1027–1047. <https://doi.org/10.1101/gad.14.9.1027>
- Porath, J. (1992). Immobilized metal ion affinity chromatography. *Protein Expression and Purification*, 3(4), 263–281. [https://doi.org/10.1016/1046-5928\(92\)90001-D](https://doi.org/10.1016/1046-5928(92)90001-D)
- Porro, D., Sauer, M., Branduardi, P., & Mattanovich, D. (2005). Recombinant protein production in yeasts. *Molecular Biotechnology*, 31(3), 245–259. <https://doi.org/10.1385/MB:31:3:245>
- Rizzuto, R., Brini, M., Pizzo, P., Murgia, M., & Pozzan, T. (1995). Chimeric green fluorescent protein as a tool for visualizing subcellular organelles in living cells. *Current Biology*, 5(6), 635–642. [https://doi.org/10.1016/S0960-9822\(95\)00128-X](https://doi.org/10.1016/S0960-9822(95)00128-X)
- Roberts, M. A. J. (2019). Recombinant DNA technology and DNA sequencing. *Essays in Biochemistry*, 63(4), 457–468. <https://doi.org/10.1042/EBC20180039>
- Rosano, G. L., & Ceccarelli, E. A. (2014). Recombinant protein expression in *Escherichia coli*: Advances and challenges. *Frontiers in Microbiology*, 5, 172. <https://www.frontiersin.org/articles/10.3389/fmicb.2014.00172>
- Rudenko, O., & Barnes, A. C. (2018). Gibson assembly facilitates bacterial allelic exchange mutagenesis. *Journal of Microbiological Methods*, 144, 157–163. <https://doi.org/10.1016/j.mimet.2017.11.023>
- Salomonsson, E., Mihalko, L. A., Luker, K. E., Verkhusha, V. V., & Luker, G. D. (2012). Cell-based and *in vivo* spectral analysis of fluorescent proteins for multiphoton microscopy. *Journal of Biomedical Optics*, 17(9),

096001. <https://doi.org/10.1117/1.JBO.17.9.096001>
- Sands, B., & Brent, R. (2016). Overview of post Cohen-Boyer methods for single segment cloning and for multisegment DNA assembly. *Current Protocols in Molecular Biology*, 113(1), 3.26.1–3.26.20. <https://doi.org/10.1002/0471142727.mb0326s113>
- Schleif, R. (2000). Regulation of the L-arabinose operon of *Escherichia coli*. *Trends in Genetics*, 16(12), 559–565. [https://doi.org/10.1016/S0168-9525\(00\)02153-3](https://doi.org/10.1016/S0168-9525(00)02153-3)
- Schleif, R., Hess, W., Finkelstein, S., & Ellis, D. (1973). Induction kinetics of the L-arabinose operon of *Escherichia coli*. *Journal of Bacteriology*, 115(1), 9–14. <https://doi.org/10.1128/jb.115.1.9-14.1973>
- Schwarze, S. R., Hruska, K. A., Dowdy, S. F., Schwarze, S. R., Hruska, K. A., & Dowdy, S. F. (2000). Protein transduction: Unrestricted delivery into all cells? *Trends in Cell Biology*, 10(7), 290–295. [https://doi.org/10.1016/S0962-8924\(00\)01771-2](https://doi.org/10.1016/S0962-8924(00)01771-2)
- Shen, Y., Chen, Y., Wu, J., Shaner, N. C., & Campbell, R. E. (2017). Engineering of mCherry variants with long Stokes shift, red-shifted fluorescence, and low cytotoxicity. *PLoS ONE*, 12(2), e0171257. <https://doi.org/10.1371/journal.pone.0171257>
- Shokolenko, I. N., Alexeyev, M. F., LeDoux, S. P., & Wilson, G. L. (2005). TAT-mediated protein transduction and targeted delivery of fusion proteins into mitochondria of breast cancer cells. *DNA Repair*, 4(4), 511–518. <https://doi.org/10.1016/j.dnarep.2004.11.009>
- Sørensen, H. P., & Mortensen, K. K. (2005). Advanced genetic strategies for recombinant protein expression in *Escherichia coli*. *Journal of Biotechnology*, 115(2), 113–128. <https://doi.org/10.1016/j.jbiotec.2004.08.004>
- Subach, F. V., Patterson, G. H., Manley, S., Gillette, J. M., Lippincott-Schwartz, J., & Verkhusha, V. V. (2009). Photoactivatable mCherry for high-resolution two-color fluorescence microscopy. *Nature Methods*, 6(2), 153–159. <https://doi.org/10.1038/nmeth.1298>
- Sun, X., Chiu, J.-F., & He, Q.-Y. (2005). Application of immobilized metal affinity chromatography in proteomics. *Expert Review of Proteomics*, 2(5), 649–657. <https://doi.org/10.1586/14789450.2.5.649>
- Sunley, K., & Butler, M. (2010). Strategies for the enhancement of recombinant protein production from mammalian cells by growth arrest. *Biotechnology Advances*, 28(3), 385–394. <https://doi.org/10.1016/j.biotechadv.2010.02.003>
- Thomas, S., Maynard, N. D., & Gill, J. (2015). DNA library construction using Gibson assembly®. *Nature Methods*, 12(11), i–ii. <https://doi.org/10.1038/nmeth.f.384>
- Wadia, J. S., & Dowdy, S. F. (2002). Protein transduction technology. *Current Opinion in Biotechnology*, 13(1), 52–56. [https://doi.org/10.1016/S0958-1669\(02\)00284-7](https://doi.org/10.1016/S0958-1669(02)00284-7)
- Wang, J. W., Wang, A., Li, K., Wang, B., Jin, S., Reiser, M., & Lockey, R. F. (2015). CRISPR/Cas9 nuclease cleavage combined with Gibson assembly for seamless cloning. *Biotechniques*, 58(4), 161–170. <https://doi.org/10.2144/000114261>
- Wang, Y., Shyy, J. Y. J., & Chien, S. (2008). Fluorescence proteins, live-cell imaging, and mechanobiology: Seeing is believing. *Annual Review of Biomedical Engineering*, 10(1), 1–38. <https://doi.org/10.1146/annurev.bioeng.010308.161731>
- Wiedenmann, J., Oswald, F., & Nienhaus, G. U. (2009). Fluorescent proteins for live cell imaging: Opportunities, limitations, and challenges. *IUBMB Life*, 61(11), 1029–1042. <https://doi.org/10.1002/iub.256>
- Wurm, F. M. (2004). Production of recombinant protein therapeutics in cultivated mammalian cells. *Nature Biotechnology*, 22(11), 1393–1398. <https://doi.org/10.1038/nbt1026>
- Yin, J., Zhu, D., Zhang, Z., Wang, W., Fan, J., Men, D., Deng, J., Wei, H., Zhang, X.-E., & Cui, Z. (2013). Imaging of mRNA–protein interactions in live cells using novel mCherry tri-molecular fluorescence complementation systems. *PLoS ONE*, 8(11), e80851. <https://doi.org/10.1371/journal.pone.0080851>
- Zhang, H., Ma, Y., Gu, J., Liao, B., Li, J., Wong, J., & Jin, Y. (2012). Reprogramming of somatic cells via TAT-mediated protein transduction of recombinant factors. *Biomaterials*, 33(20), 5047–5055. <https://doi.org/10.1016/j.biomaterials.2012.03.061>